


Guide to Identification and Ecology of New Zealand Subfossil Chironomids Found in Lake Sediment



Ann C. Dieffenbacher-Krall^{1,2}, Marcus J. Vandergoes^{1,2,3}, Craig A. Woodward⁴, and Ian K.G. Boothroyd⁵.

¹Corresponding authors: Ann C. Dieffenbacher-Krall, 112 Sawyer Research Center, University of Maine, Orono, ME 04449, USA <ann.dieffenbacher@umit.maine.edu>
Marcus J. Vandergoes, GNS Science, P.O. Box 30-368, Lower Hutt, New Zealand <m.vandergoes@gns.cri.nz>.

²Climate Change Institute, University of Maine, Orono, ME 04449, USA

³GNS Science, Lower Hutt, New Zealand

⁴Center for Chronology, Environment and Climate, Queens University, Belfast, Ireland

⁵Golder Associates (NZ) Ltd., P.O. Box 33-849, Takapuna, Auckland, New Zealand

Last updated 06/10/2008

Suggested citation:

Dieffenbacher-Krall, A.C., M.J. Vandergoes, C.A. Woodward, and I.K.G. Boothroyd. 2008.0610. Guide to Identification and Ecology of New Zealand Subfossil Chironomids Found in Lake Sediment. Climate Change Institute, University of Maine, Orono, ME <<http://www.climatechange.umaine.edu/Research/facilities/perl/nzguide.html>>.

Contact Ann Dieffenbacher-Krall to request this guide on CD.
Comments on this guide will be most welcomed.

Drawings by Dieffenbacher-Krall. Photographs by Dieffenbacher-Krall, Vandergoes, and Woodward.

This taxonomic guide is a result of work to build New Zealand subfossil chironomid training sets to enable quantitative environmental reconstructions from chironomid larvae head capsules (Dieffenbacher-Krall et al. 2007, Woodward and Shulmeister 2006). We offer this guide as a taxonomic reference for anyone working with subfossil chironomids from New Zealand or other Southern Hemisphere locations, particularly in the fields of ecological and environmental analysis and reconstruction. The guide will be updated periodically as our research progresses.

We have made this subfossil taxonomy as consistent with the current full organism taxonomy for New Zealand as possible. We have opted to use a numbering system for unnamed Orthoclaadiinae rather than the lettering system of Boothroyd (unpub.) because numbers are unlimited. Synonymy with Boothroyd's lettered Orthoclaadiinae is noted on individual pages. In many cases, the name given to a subfossil taxon was determined by examination of Boothroyd's mounts of modern larvae. We have been conservative in naming the taxa. For some New Zealand taxa there is only one species known from full organism studies. We have not used species names unless subfossil head capsules typically contain a feature distinguishing the species from other species within the genera, whether other species are found in New Zealand or not. We have not attempted to synchronize this taxonomy with the descriptions and photographs provided by Boubée (1983) and Schakau (1993) because we are not able to examine the original material. We prefer to avoid misclassification that might result from relying solely on verbal descriptions or single-view photographs provided in their theses, which may not show features critical to diagnosis of some taxa.

At this point in time, this guide is based exclusively on South Island material, although many of the types can certainly also be found on North Island, or smaller New Zealand islands. We currently recognize 60 subfossil types, of which 22 are Chironominae (5 of which are *Tanytarsus funebris* types), 2 are Diamesinae, 30 are Orthoclaadiinae, 1 is Podonominae, and 5 are Tanypodinae

Numbered Orthoclaadiinae:

In an effort to standardize subfossil chironomid taxonomy across at least a portion of the Southern Hemisphere, we have coordinated our unnamed Orthoclaadiinae numbering with Julietta Massaferro (Universidad de Bariloche, Argentina), who is currently developing a similar guide for South America. Missing numbers represent either taxa found in South America but not in New Zealand, or taxa for which we were finally able to assign a name. Retired numbers are not reused to avoid confusion.

Lakes:

Most numbered lakes listed in this document are described in Dieffenbacher-Krall et al. (2007). A few were not included in that article and information about these additional sites can be requested from Dieffenbacher-Krall. Named lakes are described in Woodward and Shulmeister (2006). For chironomid taxa found only in late-glacial age sediment, we have named the lakes. Descriptions of late-glacial chironomid studies will be forthcoming over the next several years. "Reference" describes slides made with

one head capsule per slide for the purpose of developing the taxonomy (Vandergoes and Dieffenbacher-Krall, unpub.).

We are currently working together to establish taxonomic consistency between the two inference model studies (Dieffenbacher-Krall et al. 2007, Woodward and Shulmeister 2006) based on this guide. The training sets from the two studies will then be combined. Woodward and Shulmeister (2006) lakes noted in this taxonomic guide include only those which we (Woodward, Vandergoes, and Dieffenbacher-Krall) have recounted using this standardized taxonomy. We have completed recounting 15 of the Woodward and Shulmeister (2006) sites and will update this guide as we complete more. Some of the Woodward and Shulmeister (2006) lakes were deemed inappropriate for the combined training set (e.g., too big, too much obvious human impact) and were excluded. Recounted lakes to date are Iron, Sylvester, Little Sylvester, Rainbow Skifield W, Rainbow Skifield E, Sedgemere, Princess Bath, Emma, Sylvan, Harris, Mackenzie, Gertrude Saddle/Black, Howden, Sugarbowl Tarn, and Alta.

Ecological information:

We have provided GLM models (using Canoco) for mean summer temperature (SmT), log₁₀ chlorophyll a (Chla), log₁₀ conductivity (CON), and log₁₀ percentage sediment organic content (ORG; based on 550 °C, 2 hour, loss on ignition) for all taxa (using square root of percentages) for which there is a significant (p<0.005) relationship, including only lakes from Dieffenbacher-Krall et al. (2007). When we have completed combining the two training sets, we will repeat the analyses including Woodward and Shulmeister (2006) lakes and update these pages. Methods were described by Dieffenbacher-Krall et al. (2007). Optimum and tolerance is indicated if calculated. We also generated HOF models (Huisman et al. 1993) for SmT for taxa found in a minimum of 10 sites (<http://cc.oulu.fi/~jarioksa/>; Oksanen and Minchin 2002). A Roman numeral in the corner of the graph indicates the HOF model type. For those taxa for which we have sufficient data, we provide a verbal description of subfossil ecology based solely on the sites described in Dieffenbacher-Krall et al. (2007). Subfossil chironomid ecology in lakes that are different from those included in that data set, such as North Island, highly eutrophic or disturbed, less than one meter deep, or large, may be quite different. Additional subfossil ecological information observed by Woodward and Shulmeister (2006) is noted where provided.

The data for the two training sets was collected using the same methods, but the suite of lakes included in each study is somewhat different. The altitudinal and temperature ranges of the two sets of lakes overlap, but the Dieffenbacher-Krall et al. (2007) set extends to a higher altitude and lower mean February temperatures than the Woodward and Shulmeister (2006) lake set. The latter has broader ranges of chlorophyll a and conductivity than the former (Table 1). Both studies avoided lakes with obvious human impacts.

Note well: *Subfossil head ecology and full larvae ecology are not the same.* Issues of taphonomy, how larvae head capsules are deposited and preserve in sediment,

necessitate modern studies based on the subfossil assemblage from sediment taken from the deepest part of lakes. Whole organism studies of the larval, adult, or pupal life phases may be biased with respect to season of sampling or sampling location within a lake. The preservable subfossil head capsules are extremely small and light weight and the remains of both littoral and profundal species become incorporated in deep water sediments (Iovino 1975, Wiederholm 1979, Brodin 1982, Boubée 1983, Schmäh 1993). The top half-centimeter of sediment contains subfossil material deposited over a number of years, whereas full organism sampling generally includes only those individuals living at the time of sampling. Typically, a far greater number of taxa are found by subfossil sampling than by full organism sampling.

Table 1. Range of values for various environmental variables for lakes included in Woodward and Schulmeister (2006) and Dieffenbacher-Krall et al. (2007) studies.

	# sites	Altitude m a.s.l min-max	lake depth m min-max	# taxa/lake min-max mean		SmT °C	Mean Feb. temp °C	Chla mg/m ³	CON µS/cm
Woodward and Schulmeister 2006	46	20-1882	0.74-40	1-21	10.5	n/a	8.2-18.1	0.15- 181	6.9- 1180
Dieffenbacher- Krall et al. 2007	61	120-2070	1-37	3-20	11.6	6.4-15.2	7.4-15.8	<0.1-5.9	3-96

We have provided ecological information gleaned from other sources. Note, this information is based on full organism studies unless otherwise noted. Extreme caution must be exercised when using this information to interpret subfossil assemblages.

Although researchers who study New Zealand chironomids at the organism level have been extremely productive, the size of the country and the great variety of habitat types relative to the number of researchers leave much room for continued research. Research about chironomid ecology remains in its infancy.

Definitions/Abbreviations:

cephalic setation - pattern of spots on Tanypodinae head capsules formed by pores and hair bases

Chla – chlorophyll a

CON - conductivity

Laterals - side teeth

Median - central tooth or teeth

Mentum - full set of teeth

NI – North Island

ORG – organic content (% loss-on-ignition, 550 °C, 2 hours)

SI – South Island

SmT – mean summer temperature (January, February, March)

VMP - ventromental plate

POP – post occipital plate

Additional taxa that are apt to occur in New Zealand lake sediment:

We have included only those taxa which we found represented in the subfossil assemblage. Additional taxa reported from New Zealand full organism studies that might be found as subfossil heads follow. We would greatly appreciate notification if other researchers find these or other taxa not included in this guide.

Chironominae Tanytarsini *Stempellina* (Boothroyd unpub., Stark and Winterbourn 2006): Trifid (notched) median, 6 laterals. 1st lateral smaller than 2nd. VMP striated with serrated edge. Antennae pedestals have multi-branched spurs. Known from springs and seepages in Waikato region. Found by Schakau (1993) in Lake Monowai.

Diamesinae *Heptagyini* (Stark and Winterbourn 2000): Head capsules tentatively referred to as *Heptagyini* sp. I were found by Schakau (1993) in Lady Lake. Schakau described them as having double median teeth and 5 laterals, VMP present, no beard.

Diamesinae *Lobodiamesa* (Boothroyd unpub., Boothroyd and Forsyth 2007, Stark and Winterbourn 2000): Single, broad, smooth, rounded median, 7 laterals. One species, *L. campbelli*, found occasionally in small mountain streams. A second species known from Waikato.

Orthoclaadiinae *Camptocladus* (Boothroyd unpub., Stark and Winterbourn 2000, Boothroyd and Forsyth 2007): Single, broad, rounded median, about same height as 1st lateral. 4 laterals. One species, *C. stercorarius*, is common, widespread, terrestrial, found in cattle dung.

Orthoclaadiinae *Cricotopus hollyfordensis* (Boothroyd 2002): Single median, 6 laterals. Median tooth rounded, more than 1.5 times wider and less than 1.5 times taller than 1st lateral.

Orthoclaadiinae *Cricotopus vincenti* (Boothroyd 2002): Single median, 6 laterals. Median slightly flattened. Laterals 1 and 2 partially fused with median.

Orthoclaadiinae D “Pear” (Boothroyd unpub.): Double (bifid) median, 5 laterals. Mentum sharply sloped.

Orthoclaadiinae F “Coastal” (Boothroyd unpub.): Single median, 6 laterals. 3rd laterals taller than 1 and 2. Peg-like seta subdentalis on mandibles.

Orthoclaadiinae H “Chapel” (Boothroyd unpub.): Single, pointed median, 5 laterals. 1st lateral is very short, 2nd lateral as tall as median and broader than other laterals. Laterals 3-5 descend in size and are thinner than 2. Known from Coromandel Peninsula.

Orthoclaadiinae I (Boothroyd unpub.): Double (bifid) median, 5 laterals. Laterals 3-5 small and clustered.

Orthoclaadiinae *Stictocladus* (Boothroyd unpub., Boothroyd and Forsyth 2007): Single, broad, rounded median is much taller than laterals. 4 laterals, 1st lateral taller than other 3. Laterals 2-4 are clustered together. Two species, *S. lacuniferus* and *S. pictus*, found throughout New Zealand, generally in small mountain and hill country streams, occasionally in lowland streams.

Orthoclaadiinae *Tonnoirocladius commensalis* (Cranston 2007): This is the "*Dactylocladius*" *nomen dubium* (Boothroyd unpub.): Single, broad, rounded median, 6 laterals. 1st lateral nearly as tall as median. Laterals 4-6 are on a much lower plane than median and laterals 1-3. Bearded. Found throughout New Zealand, commensal with Blephariceridae larvae, usually in cold, swift, mountain streams.

Podonominae *Podochlus* (Boothroyd unpub., Stark and Winterbourn 2006): Bifid median, 8 or 9 laterals. 3rd lateral is taller than 1 and 2. 3rd inner tooth of mandible is longer than any other mandible tooth. Four species recorded from New Zealand, but larvae unknown, probably inhabiting swift mountain streams. Larvae fitting this description found in glacial streams in Westland.

Podonominae *Podonomus* (Boothroyd unpub., Stark and Winterbourn 2006): Single median, 8 laterals. Median is about the same size as laterals 1 and 2. Lateral 3 is larger than 1 and 2. 3rd inner tooth of the mandible is longer than the apical tooth. Seven species recorded for New Zealand but larvae unknown. Probably inhabit swift mountain streams. Reported from a water-filled leaf axil of New Zealand flax.

Podonominae *Zelandochlus* (Boothroyd unpub., Dumbleton 1973, Boothroyd and Cranston 1999): Single median is taller than 1st lateral. 9 laterals. Laterals 1 and 2 roughly equal in size, lateral 3 is taller. 1st inner tooth of mandible is larger than apical tooth. One species, *Z. latipalpis*, the "ice worm", known only from the surfaces of Fox and Franz Joseph Glaciers.

Tanypodinae Macropelopini *Gressitius* (Boothroyd unpub.) See Tanypodinae General Comments.

Tanypodinae Pentaneurini *Larsia* (Boothroyd unpub.): Might be our Pentaneurini 1.

Tanypodinae Pentaneurini Pentaneurini sp. A (Boothroyd unpub.): 4 toothed ligula with all teeth roughly equal in height, bifid paraligula. About 8 teeth on pecten hypopharyngis. Found upper North Island.

Tanypodinae *Zavreliymia* (Boothroyd unpub., Boothroyd and Forsyth 2007, Stark and Winterbourn 2006): Bifid paraligula with small basal tooth. 5 toothed ligula with central

tooth being the shortest and outermost teeth angled outward. One species, *Z. harrisi*, found in streams and lakes.

Acknowledgements:

This project was funded in part by the Comer Science and Education Foundation, the National Science Foundation (ATM-0502427), the National Oceanic and Atmospheric Administration, the New Zealand Marsden Fund (contract UOC301), the New Zealand Foundation for Research Science and Technology (GNSX0501), and the Mason Scientific and Technical Trust (Department of Geological Sciences, University of Canterbury). We gratefully acknowledge taxonomic consultations with Steve Brooks, Julietta Massafiero, and participants in the Midge Dead Heads meetings (University of Fredericton, New Brunswick, May 2004, hosted by Les Cwynar; University of Okanagan, British Columbia, June 2005, hosted by Ian Walker). We thank George Denton for his role in stimulating our (Dieffenbacher-Krall and Vandergoes) work with NZ subfossil chironomids, our student chironomid extraction technicians at the University of Maine for their many hours of dedicated service, Andrea Nurse for co-supervising them, and Nicholas Dieffenbacher-Krall for stimulating discussions about chironomid life histories.

REFERENCES

- Anderson, N.H. 1982. A survey of aquatic insects associated with wood debris in New Zealand streams. *Mauri Ora* 10: 21-33.
- Boothroyd, I.K.G. 1987. Taxonomic composition and life cycles of Chironomidae (Diptera) in a northern New Zealand stream. *Entomologica scandinavia Supplements* 29: 15-22.
- Boothroyd, I.K.G. 1988. Temporal and diel emergence of Chironomidae (Diptera: Insecta) from a New Zealand stream. *Internationale Vereinigung fur Theoretische and Angewandte Limnologie* 23: 1399-1404.
- Boothroyd, I.K.G. 1989. First record of *Cricotopus aucklandensis* Sublette and Wirth (Diptera: Chironomidae) from mainland New Zealand, with a description of pupa and larva. *New Zealand Natural Sciences* 16: 51-56.
- Boothroyd, I.K.G. 1990. Additions to the Chironomidae (Diptera: Insecta) of New Zealand: *Cricotopus* van der Wulp species from a North Island stream. *New Zealand Journal of Zoology* 17: 419-436.
- Boothroyd, I.K.G. 1994. Description of *Naonella* n.gen. (Diptera: Chironomidae: Orthoclaadiinae). *New Zealand Journal of Zoology* 21: 309-315.
- Boothroyd, I.K.G. 1999a. Description of *Kaniwhaniwhanus* gen. n. (Diptera: Chironomidae: Orthoclaadiinae) from New Zealand. *New Zealand Journal of Marine and Freshwater Research* 33: 341-350.
- Boothroyd, I.K.G. 1999b. Size distribution of *Maoridiamesa stouti* Brundin (Diptera: Chironomidae): a macicolous chironomid from New Zealand. *Journal of the Kansas Entomological Society* 71 (4): 484-489.
- Boothroyd, I.K.G. 2000. Emergence patterns of Chironomidae (Diptera: Insecta) from a

- New Zealand stream. In: O. Hoffrichter, ed. Late 20th Century Research on Chironomidae: an Anthology from the 13th International Symposium on Chironomidae, Shaker Verlag, Aachen, pp. 363-374.
- Boothroyd, I.K.G. 2002. *Cricotopus* and *Paratrichocladus* (Chironomidae: Insecta) in New Zealand, with description of *C. hollyfordensis* n. sp., and redescriptions of adult and immature stages of *C. zealandicus* and *P. pluriserialis*. New Zealand Journal of Marine and Freshwater Research 36: 775-788.
- Boothroyd, I.K.G. 2004. A new species of *Naonella* Boothroyd (Chironomidae: Orthoclaadiinae) from New Zealand. New Zealand Entomologist 27: 11-15.
- Boothroyd, I.K.G., P.S. Cranston. 1995. Two Orthoclaadiinae (Chironomidae) genera common to New Zealand and Australia: *Pirara* n. gen. and *Eukiefferiella* Thienemann. In: P.S. Cranston, ed. Chironomids: From Genes to Ecosystems, CSIRO, Australia, pp. 389-408.
- Boothroyd, I., P. Cranston. 1999. The "ice worm" – the immature stages, phylogeny, and biology of the glacier midge *Zelandochlus* (Diptera: Chironomidae). Aquatic Insects 21: 303-316.
- Boothroyd, I., D. Forsyth. 2007. 1126 download (updated periodically). Checklist of New Zealand Chironomidae (Diptera). National Centre for Aquatic Biodiversity and Biosecurity. <www.niwa.co.nz/_data/assets/pdf_file/0020/432471/chirolist.pdf>
- Boubee, J.A.T. 1983. Past and present benthic fauna of Lake Maratoto with special reference to the Chironomidae. Thesis (Ph.D.), University of Waikato.
- Brodin, Y.-W. 1982. Paleoeological studies of recent development of the Lake Väjösjön. IV. Interpretation of the eutrophication process through the analysis of subfossil chironomids. Archiv für Hydrobiologie 92: 313-326.
- Brooks, S.J., P.G. Langdon, O. Heiri. 2007. The identification and use of Palaeoartic Chironomidae larvae in palaeoecology. QRA Technical Guide No. 10, Quaternary Research Association, London.
- Collier, K.J. 1993. Flow preferences of larval Chironomidae (Diptera) in Tongariro River, New Zealand. New Zealand Journal of Marine and Freshwater Research 27: 219-226.
- Collier, K.J., R.J. Wilcock, A.S. Meredith. 1998. Influence of substrate type and physico-chemical conditions on macroinvertebrate fauna and biotic indices of some lowland Waikato, New Zealand, streams. New Zealand Journal of Marine and Freshwater Research 32: 1-19.
- Cranston, P.S. 2004. Insecta: Diptera, Chironomidae. In: C.M. Yule, Y.H. Sen, eds. Freshwater Invertebrates of the Malaysian Region, Academy of Sciences, Malaysia, pp. 711-735.
- Cranston, P.S. 2007. The identity of *Dactylocladius commensalis* (Diptera: Chironomidae) revealed. Aquatic Insects 29 (2): 103-114.
- Dieffenbacher-Krall, A.C., M.J. Vandergoes, G.H. Denton. 2007. An inference model for mean summer air temperatures in the Southern Alps, New Zealand, using subfossil chironomids. Quaternary Science Reviews 26: 2487-2504.
- Dumbleton, L.J. 1973. An ice chironomid. New Zealand Entomologist 4 (2): 38-46.
- Eggermont, H., D. Verschuren. 2004a. Sub-fossil Chironomidae from East Africa. 1. Tanypodinae and Orthoclaadiinae. Journal of Paleolimnology 32: 383-412.

- Eggermont, H., D. Verschuren. 2004b. Sub-fossil Chironomidae from East Africa. 2. Chironominae (Chironomini and Tanytarsini). *Journal of Paleolimnology* 32: 413-455.
- Forsyth, D.J. 1971. Some New Zealand Chironomidae (Diptera). *Journal of the Royal Society of New Zealand* 1 (2): 113-144.
- Forsyth, D.J. 1975. Description of *Kiefferulus opalensis* n. sp. (Diptera: Chironomidae). *New Zealand Journal of Zoology* 2 (2): 215-218.
- Forsyth, D.J. 1979. Life stages and taxonomic affinities of *Xenochironomus canterburyensis* (Chironomidae: Diptera). *New Zealand Journal of Zoology* 6: 467-472.
- Forsyth, D.J. 1983. The distribution of chironomids in geothermal waters in New Zealand. *Memoirs of the American Entomological Society* 34: 115-123.
- Forsyth, D.J., I.D. McCallum. 1978. *Xenochironomus canterburyensis* (Diptera: Chironomidae), a commensal of *Hyridella menziesi* (Lamellibranchia) in Lake Taupo: features of pre-adult life history. *New Zealand Journal of Zoology* 5: 795-800.
- Graham, A.A. 1976. Ecology and production of *Chironomus zealandicus* in Lake Hayes. Thesis (M.S.), Otago University.
- Huisman, J., H. Olf, and L.F.M. Fresco. 1993. A hierarchical set of models for species response analysis. *Journal of Vegetation Science* 4: 37-46.
- Iovino, A.J. 1975. Extant chironomid larval populations and the representativeness and nature of their remains in lake sediments. Thesis (Ph.D.), Indiana University.
- Martin, J., D. Forsyth. 2007. New Zealand *Chironomus* species: larval types and karyotypes. <www.genetics.unimelb.edu.au/Martin/NZChiron.html>.
- Milner, A.M., R.C. Taylor, M.J. Winterbourn. 2001. Longitudinal distribution of macroinvertebrates in two glacier-fed New Zealand rivers. *Freshwater Biology* 46: 1765-1775.
- Oksanen, J., P.R. Minchin. 2002. Continuum theory revisited: what shape are species responses along ecological gradients? *Ecological Modeling* 157: 119-129.
- Pinder, L.C.V. 1995. Biology of the eggs and first-instar larvae. In: P.D. Armitage, P.S. Cranston, L.C.V. Pinder, eds. *The Chironomidae: Biology and Ecology of Non-Biting Midges*. Chapman & Hall, London, pp. 87-106.
- Rieradevall, M., S.J. Brooks. 2000. An identification guide to subfossil Tanypodinae larvae (Insecta: Diptera, Chironomidae) based on cephalic setation. *Journal of Paleolimnology* 23: 81-99.
- Schakau, B.L. 1993. Palaeolimnological studies on sediments from Lake Grasmere, South, New Zealand, with special reference to the Chironomidae (Diptera). Thesis (Ph.D.), University of Canterbury.
- Schmäh, A. 1993. Variation among fossil chironomid assemblages in surficial sediments of Bodensee-Untersee (SW-Germany): implications for paleolimnological interpretation. *Journal of Paleolimnology* 9: 99-108.
- Stark, J.D., M.J. Winterbourn. 2000. Chironomidae (nonbiting midges). In: M.J. Winterbourn, K.L.D. Gregson, C.H. Dolphin (eds.). *Guide to the Aquatic Insects of New Zealand*. *Bulletin of the Entomological Society of New Zealand* 13: 73-82.
- Stark, J.D., M.J. Winterbourn. 2006. Chironomidae (nonbiting midges). In: M.J.

- Winterbourn, K.L.D. Gregson, C.H. Dolphin. Guide to the Aquatic Insects of New Zealand. Bulletin of the Entomological Society of New Zealand 14: 76-84.
- Wiederholm, T. 1979. Chironomid remains in recent sediments of Lake Washington. Northwest Science 53: 251-256.
- Woodward, C.A., J. Shulmeister. 2006. New Zealand chironomids as proxies for human-induced and natural environmental change: transfer functions for temperature and lake production (chlorophyll a). Journal of Paleolimnology 36: 407-429.

Checklist of New Zealand subfossil chironomid types

	Subfamily	Tribe	Genus/species or type
1	Chironominae	Chironomini	Chironomini 1
2			Chironomini early instar
3			<i>Chironomus</i> (2 pp)
4			<i>Cladopelma</i>
5			<i>Cryptochironomus</i>
6			<i>Harrisius</i>
7			<i>Kiefferulus</i>
8			<i>Lauterborniella</i>
9			<i>Microtendipes</i>
10			<i>Parachironomus</i>
11			<i>Paucispinigera</i>
12			<i>Polypedilum</i>
13			<i>Xenochironomus</i>
14	Chironominae	Pseudochironomini	<i>Riethia/Pseudochironomini</i>
15	Chironominae	Tanytarsini	<i>Corynocera</i>
16			<i>Paratanytarsus</i>
			<i>Tanytarsus funebris</i> types (4 pp)
17			Type A short
18			Type A long
19			Type B long
20			Type C short
21			Type C long
22			<i>Tanytarsus vespertinus</i>
23	Diamesinae		Diamesinae early instar
24			<i>Maoridiamesa</i>
25	Orthoclaadiinae		<i>Corynoneura</i>
26			<i>Cricotopus aucklandensis</i>
27			<i>Cricotopus planus</i>
28			<i>Cricotopus zealandicus</i>
29			<i>Eukiefferiella</i>
30			<i>Gymnometriocnemus</i> type
31			<i>Hevelius</i>
32			<i>Kaniwhaniwhanus</i>
33			<i>Limnophyes/Paralimnophyes</i>
34			<i>Naonella forsythi</i>

35			<i>Naonella kimihia</i> (2 pp)
36			<i>Naonella</i> type 305
37			<i>Naonella</i> type 419
38			nr <i>Kaniwhaniwhanus</i>
39			Ortho 1b
40			Ortho 3
41			Ortho 4
42			Ortho 5
43			Ortho 7
44			Ortho 8
45			Ortho 9
46			Ortho 12
47			Ortho 15
48			Ortho 16
49			Ortho 19
50			<i>Parakiefferiella</i>
51			<i>Paratrichocladius</i>
52			<i>Pirara/Orthocladiinae</i> E
53			<i>Smittia</i>
54			SO4
55	Podonominae		<i>Parochlus</i>
	Tanypodinae	General comments	(1 p)
56		Macropelopini	Macropelopini type 1
57		Macropelopini	Macropelopini type 2
58		Macropelopini	Macropelopini type 3
59		Pentaneurini	<i>Ablabesmyia</i>
60		Pentaneurini	Pentaneurini 1
	Non-Chironomid	Ceratopogonidae	
		Simuliidae	
		Culicidae	